ST Large Synoptic Survey Telescope

Cosmology with Photometric Baryon Acoustic Oscillation Measurements

H. Zhan (UCD), L. Knox (UCD), J.A. Tyson (UCD) and the LSST Large-Scale Structure Science Collaboration

LSST will obtain photometric redshifts for 4 billion galaxies with the distribution peaking around z = 1 and approximately 10% of the galaxies at z > 2.5. It will achieve percent level precision on the angular diameter distance at 11 redshifts logarithmically spaced between z = 0.3 to 3.6 with a CMB-calibrated standard ruler – baryon acoustic oscillations (BAO) in the galaxy (and matter) power spectrum. By themselves, LSST BAO will provide weaker constraints on the dark energy equation of state parameters, w_0 and w_a , than LSST weak lensing (WL). However, because one can calibrate the error distribution of photometric redshifts with galaxy power spectra and determine the galaxy bias with galaxy and WL shear power spectra, a joint analysis of LSST BAO and WL will reduce the error ellipse area in the $w_0 - w_a$ plane to one sixth of that by LSST WL alone.

1. Baryon Acoustic Oscillations: A Standard Ruler

In the tightly coupled photon-plasma fluid prior to recombination, acoustic waves, supported by the photon pressure, create a characteristic scale – the sound horizon $R_{\rm S}$ in matter distribution. This results in a slight increase of correlation between density fluctuations over the scale $R_{\rm S}$ after recombination, and it appears as a series of baryon acoustic oscillations (BAO) in the galaxy and matter power spectra. The sound horizon at recombination can be determined accurately with CMB, so that BAO becomes a very promising standard ruler for measuring angular diameter distances and Hubble parameters (e.g. Eisenstein, Hu, & Tegmark 1988, ApJ, 504, L57).



3. Calibrating the Photo-z Parameters and Galaxy Bias



Gravitational lensing, being not affected by the galaxy bias, can help constrain the galaxy bias. The left panel shows the fractional error on the galaxy bias parameters from BAO (with *Planck* data) and from joint BAO and WL (also with *Planck* data, Zhan 2006, JCAP, 08, 008). The thin dashed line represents the external prior on the galaxy bias parameters, which is weak compared to the constraints from LSST data. Galaxy cross power spectra are sensitive to the photo-z error distribution, so that they can self-calibrate the photo-z parameters. This is demonstrated for the photo-z bias (middle panel) and rms error (right panel). In contrast, WL does not self-calibrate the photo-z error distribution because of its broad lensing kernel.



2. Angular Galaxy Power Spectra

Owing to its deep photometry and wide survey area, LSST will be able to obtain billions of galaxies with photometric redshifts (photo-zs) over a huge survey volume. This sample will allow for accurate measurements of the BAO features in the angular galaxy power spectra and place useful constraints on cosmological parameters.

The kernel of the galaxy power spectrum is given by the true-redshift distribution of galaxies binned by their photo-zs, which can have considerable overlap with each other. Hence, one can measure not only the auto power spectrum in each photo-z bin but also the cross power spectrum between two different photo-z bins. Moreover, the cross power spectra can provide a self-calibration of the photo-z error distribution, which is a crucial aspect of the complementarity between BAO and weak lensing (WL).



Left: Photo-z bins and the true-z distribution of galaxies. We assume that the photo-z bias $\delta_z = 0$ and the photo-z rms $\alpha_z=0.05(1+z)$ per galaxy. Right: The galaxy auto power spectrum of bin *i* at z-1.07 (solid line) and cross power spectrum between bin *i* and bin *j* (broken lines). The BAO features are prominent at 1 ~ several hundred. The grey area indicates the statistical error of the auto power spectrum at each multipole.

4. Distance Eigenmodes



The figure on the right presents uncorrelated w_0 and w_a constraints from 9 D_A eigenmodes (in 3 groups) of LSST WL and BAO. Weak priors on cosmological parameters and $\sigma_p(w_0)=1$ & $\sigma_q(w_0)=2$ are applied. The growth parameters and the galaxy bias parameters are marginalized. The solid contours include the contributions of all D_A modes. The BAO error ellipses are relatively narrow but in the same general direction, whereas the WL ones are oriented differently. The effect of combining all the BAO modes is mostly in reducing the area of the error ellipse without much change in the marginalized errors of w_0 & w_e separately.

5. Dark Energy Constraints

LSST is a public-private partnership. Design and de

As shown in the figure on the left, the constraints on w_0 and w_0 from LSST BAO are weaker than those from LSST WL. However, a *joint* analysis of BAO and WL data benefits from the extra information in the galaxy—shear power spectra, the calibration of the linear galaxy bias, and the calibration of photo-z parameters by the galaxy power spectra (Zhan 2006, astro-ph/0605696). It tightens the constraints considerably. This forecast assumes 10-year data from LSST and best current estimates of systematic error.

The figure on the left shows the best determined distance modes from LSST WL and BAO, separately (Zhan, Knox, & Tyson 2009, ApJ, 690, 923, arXiv:0806.0937). The WL modes are more concentrated at lower redshifts than the BAO modes, because WL measures the fluctuations at lower redshifts

measures the fluctuations at lower redshifts than the source galaxies and because the galaxy power spectra are truncated more at lower redshifts to reduce the impact of the nonlinear evolution. Although in general BAO places tighter constraints on individual D_A s than WL, the WL modes shown actually have smaller errors than the BAO modes.

