


# LSST

Large Synoptic Survey Telescope

## Mapping the Milky Way with LSST

J. S. Bullock (UC Irvine), C. M. Rockosi (UC Santa Cruz), Z. Ivezić (U. Washington), A. Saha (NOAO), the LSST Milky Way Science Collaboration

Recent results from large-area surveys have drastically altered our view of the Galaxy, revealing a complex, dynamical structure that is still being shaped by the infall of smaller systems. The multicolor, multi-epoch photometric map created by the LSST will provide an unprecedented means to extend our exploration of the Galaxy's structure, star formation, chemical enrichment, and accretion history on a panoramic scale. Strategic time and space sampling of each field over ten years will allow variability, proper motion and parallax measurements for objects brighter than V=24, and will permit both photometric parallaxes and chemical abundance estimates for over a billion main sequence stars to 100 kpc. New dwarf galaxies in the Local Group and other nearby groups will be revealed via over-densities in resolved stars and provide important constraints on the nature of the Dwarf Satellite problem faced by CDM cosmologies.




### Structure of the Outer Milky Way:

Left: Star density multiplied by radius cubed from the stellar halo simulations of Bullock & Johnston (2005). LCDM predicts that the Milky Way should have accreted and destroyed hundreds of small dwarf galaxies in the past 10 Gyr. The residue of this process survives as structure in the outer halo. Overdensities of this kind have been revealed by SDSS star count studies, including RR Lyrae stars to a distance of 100 kpc (Ivezić et al. 2003).

- LSST will detect RR Lyrae to 400 kpc, extending the SDSS mapping area by a factor of 50. This will provide an important test of the small-scale accretion history of the Galaxy and a test of LCDM.

- RR Lyrae maps and related main sequence star counts with kinematics will allow precise measurements of the Milky Way dark halo potential.




### Structure of the Inner Milky Way:

Left: Surface brightness maps from a disk galaxy simulation by Kazantzidis et al. (2007). Faint, complex structures arise from its hierarchical merging history. Star count maps offer a unique means to constrain theories by revealing detailed features. The first generation of studies have found similar structures in the Milky Way and M31 (Newberg et al. 2002, Ferguson et al. 2002, Juric et al. 2007) and offer a glimpse into the dynamic formation history of large galaxies.

- LSST will provide photometric parallaxes for over a billion main sequence stars to 100 kpc and map the Galaxy's surroundings to high precision.

- Time series data will provide proper motions at ~10 km/s at 10 kpc (dashed circle) and chemical abundance estimates for main sequence stars. Multi-dimensional maps will produce a phase-space record of the Galaxy's formation history.



### Dwarf Satellites and the Nature of Dark Matter:


Left: Predictions for the dark subhalo velocity function in Milky-Way size halos for LCDM ( $\sigma_8=0.83$ , left panel) and WDM (right panel) from Zentner & Bullock (2003). Red squares show the cumulative dwarf satellite count prior to SDSS. The green square includes 8 new low-surface brightness dwarf spheroidals discovered using star counts in SDSS maps (e.g. Willmann et al. 2005, Belokurov et al. 2007).

- Fainter star count maps over more sky will allow LSST to reveal new dwarfs as stellar overdensities. This possibility is represented by the blue square in the figures.

- A more complete census of the Milky Way satellites will help elucidate the nature of the dwarf satellite problem facing CDM. The current count (green) is close to the WDM prediction and new discoveries may potentially rule out interesting WDM models.

### References:

- Belokurov, V. et al. 2007, astro-ph/0608448
- Bullock, J. S. & Johnston, K. V. 2005, ApJ 635, 931
- Bullock, J. S., Kravtsov, A. V., & Weinberg, D. H., 2000, ApJ, 539, 517
- Ferguson, A. M. N. et al. 2002, AJ 124, 1452
- Ivezić, Z. et al. 2003, astro-ph/030905
- Juric, M. et al. 2007, astro-ph/0510520
- Kazantzidis, S., Zentner, A. R., Bullock, J. S., & Kravtsov, A. V., 2007
- Willmann, B. et al. 2005, ApJ, 626, L85
- Zentner, A. R. & Bullock, J. S. 2003, ApJ, 598, 49



The effort to build the Large Synoptic Survey Telescope is overseen by the LSST Corporation, a non-profit 501(c)(3) corporation formed in 2003, with headquarters in Tucson, AZ. The LSST research and development effort is funded in part by the National Science Foundation under Scientific Program Order No. 9 (AST-0551161) through Cooperative Agreement AST-0132798. Additional funding comes from private donations, in-kind support at Department of Energy laboratories and other LSSTC Institutional Members.

Research Corporation • University of Arizona • NOAO • University of Washington • Brookhaven National Laboratory (BNL) • Harvard-Smithsonian Center for Astrophysics (CIA) • Johns Hopkins University • Las Cumbres Observatory, Inc. • Lawrence Livermore National Laboratory (LLNL) • Stanford Linear Accelerator Center (SLAC) • Stanford University Kavli Institute for Particle Astrophysics (KIPAC) • University of Pennsylvania • The Pennsylvania State University • University of California, Davis • University of California, Irvine • University of Illinois at Urbana-Champaign • Google Inc.